

Ibuprofen (Motrin or Advil) Dosing for Infants and Children

May be given every 6-8 hours; maximum of 4 times in 24 hours.

Not recommended for infants under 12 lbs or those under 6 months of age OR in people with dehydration.

Weight (pounds)	Weight (kg)	Dose (mg)	Children's Suspension (100mg/5ml)	Children's Chewables(50mg each)	Jr Strength Tablets (100mg each)	Adult (200 mg each)
11 – 16 lbs	5 – 7 kg	50 mg	2.5 ml			
16 – 22 lbs	7 – 10 kg	75 mg	3.75 ml			
22 – 28 lbs	10 – 13 kg	100 mg	5.0 ml	2 tablets	1 tablet	
28 – 33 lbs	13 – 15 kg	125 mg	6.25 ml	2.5 tablets	1.5 tablets	
33 – 44 lbs	15 – 20 kg	150 mg	7.5 ml	3 tablets	2 tablets	
44 – 88 lbs	20 – 40 kg	200 mg	10.0 ml	4 tablets	2.5 tablets	1
88 – 130 lbs	40 – 59 kg	300 mg	15.0 ml	6 tablets	3 tablets	1.5
Over 130 lbs	Over 59 kg	400 mg	20.0 ml	8 tablets	4 tablets	2

Acetaminophen (Tylenol) Dosing for Infants and Children

May be given every 4-6 hours; maximum 5 times in 24 hours.

WARNING: Infant Drops must use the dropper provided to avoid risk of OVERDOSE

Weight (pounds)	Weight (kg)	Dose (mg)	Infant Suspension (160mg/5ml)	Children's Suspension(160mg/5ml)	Children's Tablets (80mg)	Jr Strength Tablets (160mg)	Adult (325 mg)
6 – 12 lbs	3 – 5 kg	40 mg	1.25 ml	1.25 ml			
12 – 18 lbs	5 – 8 kg	80 mg	2.50 ml	2.50 ml			
18 – 24 lbs	8 – 11 kg	120 mg	3.75 ml	3.75 ml			
24 – 36 lbs	11 – 16 kg	160 mg	5.0 ml	5.0 ml	2 tablets	1 tablet	
36 – 48 lbs	16 – 22 kg	240 mg	7.5ml	7.5 ml	3 tablets	1.5 tablets	
48 – 60 lbs	22 – 27 kg	320 mg		10.0 ml	4 tablets	2 tablets	
60 – 72 lbs	27 – 33 kg	400 mg		12.5 ml	5 tablets	2.5 tablets	
72 – 95 lbs	33 – 43 kg	480 mg		15.0 ml	6 tablets	3 tablets	
95 – 150 lbs	43 – 68 kg	640 mg		20.0 ml	8 tablets	4 tablets	2
Over 150 lbs	Over 68 kg	960 mg			12 tablets	6 tablets	3